

LA-UR-16-28155

Approved for public release; distribution is unlimited.

Title: D-wave Quantum Computer as an Efficient Classical Sampler

Author(s): Chertkov, Michael
Hagberg, Aric Arild
Lokhov, Andrey
Misiakiewicz, Theodor
Misra, Sidhant
Vuffray, Marc Denis

Intended for: Distribution of slides from the ISTI D-wave rapid response call

Issued: 2016-10-26 (Draft)

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



D-wave Quantum Computer as an Efficient Classical Sampler

M. Chertkov^{1,2} (PI), A. Hagberg³, A. Lokhov^{1,2} (co-PI),
T. Misiakiewicz¹, S. Misra³, M. Vuffray²

¹Center for Nonlinear Studies

²Theoretical Division T-4

³Theoretical Division T-5

D-wave Quantum Computing Efforts Debrief

Introduction: D-wave as an efficient sampler


Theoretical and experimental evidence that D-wave can approximately sample from a Boltzmann distribution at some effective temperature

Ronnow *et al.*, Science (2014)

Amin, Phys. Rev. A (2015)


Perdomo-Ortiz *et al.*, Sci. Rep. (2016)

Benedetti *et al.*, Phys. Rev. A (2016)



Introduction: D-wave as an efficient sampler

Disadvantage for optimization turned into **advantage** for numerous applications:



- ✓ **Restricted Boltzmann Machines** (blocks for Deep Learning)
Denil & Freitas, NIPS (2011); Dumoulin *et al.*, AAAI Artificial Intelligence (2015); Benedetti *et al.*, Phys. Rev. A (2016); Amin *et al.*, "Quantum Boltzmann Machine" (2016)
- ✓ **Producing samples in hard glassy models**
Katzgraber *et al.*, Phys. Rev. X (2014 & 2015); Martin-Mayor & Hen, Sci. Rep. (2015); Venturelli *et al.*, Phys. Rev. X (2015); Zhu *et al.*, Phys. Rev. A (2016)
- ✓ **Accurate calibration** of the D-wave machine
King & McGeoch (2014) "Algorithm engineering for a quantum annealing platform"; Perdomo-Ortiz *et al.*, Sci. Rep. (2016); Raymond *et al.*, "Global warming: temperature estimation in annealers" (2016); Also example in this debrief!

Relation between input and effective Hamiltonians in D-wave

Input Hamiltonian

$$\mathcal{H} = \sum_{\langle i,j \rangle} J_{ij} \sigma_i \sigma_j + \sum_{i \in V} H_i \sigma_i$$

Effective Hamiltonian in D-wave

$$\mathcal{H}_{\text{eff}} = \sum_{\langle i,j \rangle} J'_{ij} \sigma_i \sigma_j + \sum_{i \in V} H'_i \sigma_i$$

Relation between input and effective Hamiltonians in D-wave

Input Hamiltonian

$$\mathcal{H} = \sum_{\langle i,j \rangle} J_{ij} \sigma_i \sigma_j + \sum_{i \in V} H_i \sigma_i$$

Effective Hamiltonian in D-wave

$$\mathcal{H}_{\text{eff}} = \sum_{\langle i,j \rangle} J'_{ij} \sigma_i \sigma_j + \sum_{i \in V} H'_i \sigma_i$$

Let us write $J'_{ij} = \beta(J_{ij} + \Delta J_{ij})$, $H'_i = \beta(H_i + \Delta H_i)$, where

$T = 1/\beta$: effective temperature

ΔJ_{ij} , ΔH_i : possible biases

Correspondence $\mathcal{H} \leftrightarrow \mathcal{H}_{\text{eff}}$ by solving the reconstruction problem of learning β , ΔJ_{ij} , ΔH_i from samples produced by D-wave with \mathcal{H}_{eff}

Reconstruction problem in D-wave

Given M independent **samples** (configurations), **reconstruct** \mathcal{H}_{eff}

	$k = 1$	$k = 2$...	$k = M$
σ_1	+1	-1	...	+1
σ_2	-1	-1	...	-1
\vdots	\vdots	\vdots	...	\vdots
σ_N	+1	+1	...	-1

Task known as **Inverse Ising Problem**. The optimal algorithm for solving this task is the **LANL-developed “Screening method”**

Vuffray, Misra, Lohkov, Chertkov, NIPS (2016)
Lohkov, Vuffray, Misra, Chertkov, submitted to Nature Physics (2016)


How does Screening method work?

For each spin, **minimize** the potential $S_i(J_i, H_i)$ which **applies counter-interactions** ($P \propto e^{-\mathcal{H}}$):

$$(\hat{J}_i, \hat{H}_i) = \underset{(J_i, H_i)}{\operatorname{argmin}} \left(S_i(J_i, H_i) + \lambda \|J_i\|_1 \right)$$

$$S_i(J_i, H_i) = \langle \exp(\sum_{j \neq i} J_{ij} \sigma_i \sigma_j + H_i \sigma_i) \rangle_M$$

Vuffray, Misra, Lokhov, Chertkov, NIPS (2016)
Lokhov, Vuffray, Misra, Chertkov, submitted to Nature Physics (2016)



First outcome of this project: development of an **efficient algorithmic implementation** using advanced first-order optimization methods ($\sim N^2$ times faster, to appear on GitHub)

Effective temperature

Where does the **effective temperature** come from? Let us look at the annealing procedure with $\tau = t/t_{\text{annealing}}$:

$$\mathcal{H}(\tau) = A(\tau) \left(- \sum_{i \in V} \sigma_i^x \right) + B(\tau) \left(\sum_{\langle i,j \rangle} J_{ij} \sigma_i^z \sigma_j^z + \sum_{i \in V} H_i \sigma_i^z \right)$$

Monotonic functions A and B satisfy $A(0) \gg B(0)$ and $A(1) \ll B(1)$.

Effective temperature

Where does the **effective temperature** come from? Let us look at the annealing procedure with $\tau = t/t_{\text{annealing}}$:


$$\mathcal{H}(\tau) = A(\tau) \left(- \sum_{i \in V} \sigma_i^x \right) + B(\tau) \left(\sum_{\langle i,j \rangle} J_{ij} \sigma_i^z \sigma_j^z + \sum_{i \in V} H_i \sigma_i^z \right)$$

Monotonic functions A and B satisfy $A(0) \gg B(0)$ and $A(1) \ll B(1)$.

The “**freeze-out**” phenomenon: the evolution stops at the point τ_{freeze} :

$$T_{\text{eff}} = T_{\text{D-wave}} \frac{B(1)}{B(\tau_{\text{freeze}})}$$



Benedetti *et al.*, Phys. Rev. A (2016)
Raymond *et al.*, “Global warming: temperature estimation in annealers” (2016)



- ✓ **No unique T_{eff} :** β is the **function of the input Hamiltonian**
- ✓ “**Single qubit freeze-out**”: τ_{freeze} can **vary for different spins**

Illustration: estimating the effective temperature

Data set (from **Marcus Daniels**): embedded closed circles of $N = 22$ spins with different values of $J_{i,i+1}$ and $H_i = 0$ (diverse realizations, $t_{\text{annealing}}$, etc.). Example for $M = 7250$ and $J_{i,i+1} = -0.0625 \forall (i, i+1)$.



Refined $\{J'_{ij}, H'_i\}$. Neglecting H'_i and biases, $\beta_{\text{eff}} \approx 7$ since $\overline{J'} = -0.44$.







Illustration: estimating the effective temperature

Example for $M = 7250$ and $J_{i,i+1} = -0.4375 \forall (i, i+1)$.



Refined $\{J'_{ij}, H'_i\}$. Neglecting H'_i and biases, $\beta_{\text{eff}} \approx 4.2$ since $\overline{J'} = -1.84$.







Illustration: estimating the effective temperature

Example for $M = 7250$ and $J_{i,i+1} = -0.75 \forall (i, i+1)$.



Refined $\{J'_{ij}, H'_i\}$. Neglecting H'_i and biases, $\beta_{\text{eff}} \approx 3.72$ since $\overline{J'} = -2.79$.







Illustration: estimating the effective temperature

In the case of $J_{i,i+1} = -1.0 \forall (i, i+1)$, $M = 7250$ is insufficient: the topology can not be correctly recovered.




Dependence between J' on J :




What about biases?

Simple test: if $P(\underline{\sigma}) \propto e^{-\mathcal{H}(\underline{\sigma})/(\alpha T)}$, then $\alpha^2 T \frac{\partial}{\partial \alpha} \langle H \rangle = \langle H^2 \rangle - \langle H \rangle^2$

Checkerboard pattern with magnetic fields



Checkerboard pattern without magnetic fields



Example found by **Carleton Coffrin**, [see next talk!](#)

Illustration: detecting and correcting biases

Example of the input $\mathcal{H} = 0$ over the entire Chimera graph

Although D-wave comes with a software for correcting biases, they are **still present and persist**. Example from the Burnaby machine on Sep 15:






Illustration: detecting and correcting biases

September 15



October 4





Illustration: detecting and correcting biases







Illustration: detecting and correcting biases











Illustration: detecting and correcting biases

Corrections: inputting $\mathcal{H} = -\frac{1}{\beta_J} \sum_{\langle ij \rangle} J_{ij}^{(bias)} \sigma_i \sigma_j - \frac{1}{\beta_H} \sum_{i \in V} H_i^{(bias)} \sigma_i$

\mathcal{H}_{eff} before corrections

\mathcal{H}_{eff} after corrections



Symmetrized and more squeezed distributions with a single iteration

Path forward: efficient calibration of the D-wave machine

The **calibration issue** addressed in several recent papers with heuristic methods: King & McGeoch (2014); Perdomo-Ortiz *et al.*, Sci. Rep. (2016); ...

As shown in the previous examples, **we can do much better!**

✓ **Iteratively correcting the biases** for the target \mathcal{H}_T :

$$i) \frac{\mathcal{H}_T}{\beta} \longrightarrow \mathcal{H}_T + \Delta(\mathcal{H}_T)$$

$$ii) \frac{\mathcal{H}_T - \Delta(\mathcal{H}_T)}{\beta} \longrightarrow \mathcal{H}_T - \Delta(\mathcal{H}_T) + \Delta(\mathcal{H}_T - \Delta(\mathcal{H}_T))$$
$$\approx \mathcal{H}_T - \Delta'(\mathcal{H}_T)\Delta(\mathcal{H}_T)$$

$$iii) \frac{\mathcal{H}_T - \Delta(\mathcal{H}_T) + \Delta'(\mathcal{H}_T)\Delta(\mathcal{H}_T)}{\beta} \longrightarrow \dots$$

✓ **Machine learning task:** **learn the functional form** of $\Delta(\mathcal{H}_T)$ with the **linear response theory**; start directly at the point (ii)

✓ Include the **higher-order interaction** terms in the reconstruction problem to capture the effect of **inactive spins**

Acknowledgements and questions

Many thanks to Marcus Daniels for the data set used in the [first part](#) of the work, and to Carleton Coffrin for the insight and data contributions in the [second part](#)!

Questions?

